

LIST OF FIGURES

Figure 2-1.	Average concentrations of total suspended solids, biochemical oxygen demand, and oil and grease in 5-Mile Outfall effluent from reporting years 1986-87 through 2005-06.....	10
Figure 2-2.	Concentrations of arsenic, chromium, and lead in plant influent and 5-Mile Outfall effluent for reporting years 1986-87 through 2005-06.	11
Figure 2-3.	Concentrations of nickel, copper, and zinc in plant influent and 5-Mile Outfall effluent for reporting years 1986-87 through 2005-06.	12
Figure 3-1.	Historical rainfall in Los Angeles, 1877-2005 (NOAA, 2007).....	31
Figure 3-2.	Monthly rainfall amounts, 2005 and 2006.	32
Figure 3-3.	Water quality stations samples in Santa Monica Bay, 2005-2006	33
Figure 3-4.	CTD and Rosette sampler being deployed.....	34
Figure 3-5.	Microbiology inshore and shoreline stations in Santa Monica Bay, 2005-2006	35
Figure 3-6.	Calculated salinity anomaly (S_A) and computed salinity (S) values for various dilutions based on a mean salinity of 33.346 psu.....	37
Figure 3-7.	Bay-wide low salinity February 2005	41
Figure 3-8a.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in February of 2005.....	42
Figure 3-8b.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in May of 2005. Note that conductivity sensor failure caused partial data loss of the 3300, 3500 and the 3700 transects.....	43
Figure 3-8c.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in August of 2005	44
Figure 3-8d.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in November of 2005.....	45

Figure 3-8e.	CDOM plots of the near-outfall across-shelf transects for February and May Surveys	46
Figure 3-8f.	CDOM plots of the near-outfall across-shelf transects for August and November 2005 Surveys.....	47
Figure 3-9a.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in February of 2006.....	49
Figure 3-9b.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in May of 2006	50
Figure 3-9c.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in August of 2006	51
Figure 3-9d.	Temperature and Salinity 2-D plots of the 3206-4006, 3205-4005 and 3204-4004 along-shelf transects with water column depths ranging from 10-m to a maximum of 100-m for the survey conducted in November of 2006. Note that conductivity sensor failure caused data to be lost for the 3800 transect	52
Figure 3-9e.	CDOM plots of the near-outfall across-shelf transects for February and May 2006 Surveys.....	53
Figure 3-9f.	CDOM plots of the near-outfall across-shelf transects for August and November 2006 Surveys.....	54
Figure 3-10.	2005 annual geometric means and 2006 single sample results for indicator bacteria at inshore (surface and depth) stations in Santa Monica Bay	55
Figure 3-11.	Annual geometric means for indicator bacteria at shoreline stations in Santa Monica Bay, Jan 2005 – May 2005	56
Figure 3-12.	Monthly geometric means for indicator bacteria for Hyperion Treatment Plant effluent January –May 2005	57
Figure 4-1a.	Sediment chemistry station sampled in Santa Monica Bay during 2005.....	70
Figure 4-1b.	Sediment chemistry station sampled in Santa Monica Bay during 2006.....	71
Figure 4-2a.	Percentage of (Silt + Clay) in surface sediment of Santa Monica Bay in summer 2005.....	74

Figure 4-2b.	Percentage of (Silt + Clay) in surface sediment of Santa Monica Bay in summer 2006.....	74
Figure 4-3a.	Concentrations of TOC in surface sediments of Santa Monica Bay in summer 2005.....	75
Figure 4-3b.	Concentrations of TOC in surface sediments of Santa Monica Bay in summer 2006.....	75
Figure 4-4a.	Concentrations of metals Ag, As, and Pb in surface sediments of Santa Monica Bay in summer 2005.....	78
Figure 4-4b.	Concentrations of metals Ag, As, and Pb in surface sediments of Santa Monica Bay in summer 2006.....	78
Figure 4-5a.	Concentrations of metals Ni and Cr in surface sediments of Santa Monica Bay in summer 2005.	79
Figure 4-5b.	Concentrations of metals Ni and Cr in surface sediments of Santa Monica Bay in summer 2006.	79
Figure 4-6a.	Concentrations of metals Hg and Cd in surface sediments of Santa Monica Bay in summer 2005.....	80
Figure 4-6b.	Concentrations of metals Hg and Cd in surface sediments of Santa Monica Bay in summer 2006.....	80
Figure 4-7a.	Concentrations of metals Cu and Zn in surface sediments of Santa Monica Bay in summer 2005.....	81
Figure 4-7b.	Concentrations of metals Cu and Zn in surface sediments of Santa Monica Bay in summer 2006.....	81
Figure 4-8a.	Concentrations of Total PCBs and Total DDTs in surface sediments of Santa Monica Bay in summer 2005.	82
Figure 4-8b.	Concentrations of Total PCBs and Total DDTs in surface sediments of Santa Monica Bay in summer 2006.	82
Figure 4-9.	Concentrations (mg/kg) of arsenic in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	83
Figure 4-10.	Concentrations (mg/kg) of cadmium in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	84
Figure 4-11.	Concentrations (mg/kg) of chromium in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	84

Figure 4-12.	Concentrations (mg/kg) of copper in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	85
Figure 4-13.	Concentrations (mg/kg) of lead in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	85
Figure 4-14.	Concentrations (mg/kg) of mercury in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	86
Figure 4-15.	Concentrations (mg/kg) of nickel in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	87
Figure 4-16.	Concentrations (mg/kg) of silver in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	87
Figure 4-17.	Concentrations (mg/kg) of zinc in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	88
Figure 4-18.	Concentrations (mg/kg) of total DDTs in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	88
Figure 4-19.	Concentrations (mg/kg) of total PCBs in sediment at E6 (7-Mile), Z2 (5-Mile), and northern reference site (C1) from 2003 to 2006.....	89
Figure 5-1.	Benthic macrofauna stations sampled in winter 2005 and summer 2006.	98
Figure 5-2.	Benthic macrofauna stations sampled in summer 2005.	99
Figure 5-3.	Mirror tree mapping of species richness (S, left) and total abundance (N, right).	107
Figure 5-4.	Mirror tree mapping of average taxonomic distinctness (Δ^+ , left) and benthic response index (BRI, right).	108
Figure 5-5.	Parsimony analysis of endemicity (PAE) cladogram of Santa Monica Bay macrofauna stations, 2005-2006.....	110
Figure 5-6.	Mirror tree mapping of sand (left) and silt (right) fractions on PAE cladogram of Santa Monica Bay macrofauna stations, 2005-2006.....	111
Figure 5-7.	Depth mapped onto PAE cladogram of Santa Monica Bay macrofauna stations, 2005-2006.....	112
Figure 5-8.	3-D plot mirror tree mapping (depth and granulometry PCA ordination) of depth (left) and silt (right) of Santa Monica Bay macrofauna stations, 2005-2006.....	113

Figure 5-9.	Mirror tree mapping of <i>Amphiodia urtica</i> (left) and <i>Glycera macrobranchia</i> (right) on PAE cladogram of Santa Monica Bay macrofauna stations, 2005-2006.....	114
Figure 5-10.	<i>Amphiodia urtica</i> abundance layered on GIS map of Santa Monica Bay.....	115
Figure 5-11.	Parsimony analysis of endemicity (PAE) cladogram of the subset of stations sampled for priority pollutants.	116
Figure 5-12.	3-D plot mirror tree mapping (depth, granulometry, metals, and total DDT PCA ordination) of sand (left) and depth (right) of Santa Monica Bay macrofauna stations, 2005-2006.	117
Figure 5-13.	3-D plot mirror tree mapping (depth, granulometry, metals, and total DDT PCA ordination) of total DDT (left) and zinc (right) of Santa Monica Bay macrofauna stations, 2005-2006.....	118
Figure 5-14.	Parsimony analysis of endemicity (PAE) cladogram of 7-Mile Outfall station E6, 1983-2006.....	119
Figure 5-15.	3-D plot tree of PAE cladogram (macrofauna PCA ordination), black spots 1983-1987, green spots 1988-2006 after cessation of sludge discharge.	120
Figure 5-16.	Mirror tree mapping of species richness (S, left) and total abundance (N, right) on PAE cladogram of Station E6 over time.	121
Figure 5-17.	Mirror tree mapping of sand (left) and clay (right) on constellation PAE cladogram of station E6 over time.....	123
Figure 5-18.	Mirror tree mapping of BRI (left) and Delta ⁺ (right) on constellation PAE cladogram of station E6 over time.....	124
Figure 5-19.	Parsimony analysis of endemicity (PAE) cladogram of 5-Mile Outfall stations Z1 and Z2, 1983-2006.....	125
Figure 5-20.	3-D plot tree of PAE cladogram (macrofauna PCA ordination), black spots 1983-1998, green spots 1999-2006 after full secondary treatment went on line.....	129
Figure 5-21.	2-D and 3-D NMDS ordinations derived from patristic distance matrix from PAE cladogram.....	130
Figure 5-22.	Mirror tree mapping of suspended solids (left) and Shannon diversity (H', right) on PAE cladogram of Stations Z1 and Z2 over time.....	131
Figure 5-23.	Mirror tree mapping of BRI (left) and total phylogenetic diversity (Sphi ⁺ , right) on PAE cladogram of Stations Z1 and Z2 over time.....	132

Figure 5-24.	Mirror tree mapping of <i>Euphilomedes carcharodonta</i> (left) and <i>Parvilucina tenuisculpta</i> (right) on PAE cladogram of Stations Z1 and Z2 over time.	135
Figure 5-25.	Mirror tree mapping of <i>Amphiodia urtica</i> (left) and <i>Eunice americana</i> (right) on PAE cladogram of Stations Z1 and Z2 over time.	136
Figure 6-1.	Trawl stations sampled in Santa Monica Bay, winter 2005 and summer 2006.....	147
Figure 6-2.	<i>Lytechinus pictus</i>	155
Figure 6-3.	2005-2006 PAE depth-mapped cladogram.	156
Figure 6-4.	2005-2006 Q-mode NMDS with depth factor applied.	158
Figure 6-5.	2005-2006 Q-mode NMDS with spatial factor applied.	159
Figure 6-6.	1983-2006 5-Mile Outfall PAE cladogram.	160
Figure 6-7.	3-D plot tree of 5-Mile Outfall PAE cladogram (within-group PCA ordination), black spots 1983-1998, green spots 1999-2006 after full secondary treatment went on line.	161
Figure 6-8.	1983-2006 PAE trawl response index-mapped cladogram.	162
Figure 6-9.	1983-2006 PAE Sphi+-mapped cladogram.....	164
Figure 6-10.	1983-2006 PAE cladogram with <i>Citharichthys sordidus</i> abundance mapped.....	169
Figure 7-1.	Fish tissue bioaccumulation zones for both Local Seafood Safety and Local Bioaccumulation Trends Surveys in Santa Monica Bay, 2005-2006.	176
Figure 7-2.	Arsenic concentrations (mg/kg) in muscle tissues of hornyhead turbot from all three zones during 2002 through 2006 surveys. No samples were collected from Zone 5 in August 2002, and from the Nearfield Zone in August 2005....	183
Figure 7-3.	Selenium concentration (mg/kg) in muscle tissues of hornyhead turbot at three zones during 2005 and 2006 surveys. No sample was collected at the Nearfield Zone in August 2005. No selenium data available for all three zones from February 2002 to March 2005.	184
Figure 7-4.	Mercury concentrations (mg/kg) in muscle tissues of hornyhead turbot from all three zones during 2002 through 2006 surveys. No sample was collected from the Nearfield Zone in August 2005	185

Figure 7-5.	Concentrations (mg/kg) of total DDT in muscle tissues of hornyhead turbot at three zones during 2002 through 2006 surveys. No sample was collected from the Nearfield Zone in August 2005.	186
Figure 7-6.	Concentrations (mg/kg) of total PCB in muscle tissues of hornyhead turbot at three zones during 2002 through 2006 surveys. No sample was collected from the Nearfield Zone in August 2005, Zone 5 in March 2005, or Zone 4 in March 2005.	187
Figure 7-7.	Concentrations of % lipid in muscle tissues of hornyhead turbot at three zones during the 2002 through 2006 surveys. No samples at Nearfield Zone in August 2005.	188
Figure 7-8.	Arsenic concentration (mg/kg) in liver tissues of hornyhead turbot at three zones during the 2002 through 2006 surveys. No samples at Zone 5 in August 2002, and at Nearfield Zone in August 2005.	190
Figure 7-9.	Selenium concentration (mg/kg) in liver tissues of hornyhead turbot at three zones during 2005 and 2006 surveys. No sample was collected from the Nearfield Zone in August 2005. Analysis of Se was required only after the issuance of the new permit in April 2005.	191
Figure 7-10.	Mercury concentration (mg/kg) in liver tissues of hornyhead turbot at three zones during the 2002 through 2006 surveys. No sample was collected from the Nearfield Zone in August 2005.	192
Figure 7-11.	Concentration (mg/kg) of total DDT in liver tissues of hornyhead turbot from three zones during the 2002 through 2006 surveys. No sample was collected from the Nearfield Zone in August 2005.	193
Figure 7-12.	Concentration (mg/kg) of total PCB in liver tissues of hornyhead turbot from three zones during 2002 through 2006 surveys. No sample was collected from the Nearfield Zone in August 2005.	194
Figure 7-13.	Concentration of %lipid in liver tissues of hornyhead turbot from three zones during 2002 through 2006 surveys. No samples were collected from the Nearfield Zone in August 2005, or from Zone 4 (south SMB) in August 2005 and August 2006.	195
Figure 7-14.	Concentrations of arsenic, selenium, and mercury in muscle tissues of California scorpionfish at 5-Mile Outfall during the 2001 through 2005 surveys.	198
Figure 7-15.	Concentration of arsenic, selenium, and mercury in muscle tissues of vermilion rockfish at 5-Mile Outfall during the 2001 through 2005 surveys.	199