

LIST OF TABLES

Table 1-1.	Summary of the City of Los Angeles' current ocean monitoring program for Santa Monica Bay, with 1998-99 modifications and 2005 permit changes.....	3
Table 2-1.	Constituents measured in the current 5-Mile Outfall effluent monitoring program, effective May 14, 2005.....	9
Table 2-2a-1.	Annual averages, maximum and minimum concentrations of conventional constituents and other pollutants in the 5-Mile Outfall effluent (January 2005 through May 13, 2005). All concentrations are reported in µg/L unless otherwise noted.....	13
Table 2-2a-2.	Annual averages, maximum and minimum concentrations of conventional constituents and other pollutants in the 5-Mile Outfall effluent (May 14, 2005 through December 2005). All concentrations are reported in µg/L unless otherwise noted.....	15
Table 2-2b.	Annual averages, maximum and minimum concentrations of conventional constituents and other pollutants in the 5-Mile Outfall effluent (January 2006 through December 2006). All concentrations are reported in µg/L unless otherwise noted.....	17
Table 2-3.	Average <i>monthly</i> averages of total suspended solids (TSS), Biochemical Oxygen Demand (BOD), oil & grease (O&G), and settleable solids in plant influent and 5-Mile Outfall effluent during January 2005 through December 2005 and January 2006 through December 2006.....	19
Table 2-4a.	Average, maximum, and minimum values of detected priority pollutant metals and cyanide (µg /L) in plant influent and 5-Mile Outfall effluent during January 2005 through December 2005.....	20
Table 2-4b.	Average, maximum, and minimum values of detected priority pollutant metals and cyanide (µg /L) in plant influent and 5-Mile Outfall effluent during January 2006 through December 2006.....	21
Table 2-5.	Concentrations (µg/L) of key metals in 5-Mile Outfall effluent and their removal efficiency for years 1986-87 through 2005-06.	22
Table 3-1.	NPDES receiving water limitations in 2005 permit.....	30
Table 3-2.	Percent of NPDES bacteriological standards limits compliance with total coliform standards (1-5) and <i>E. coli</i> ** standards (6-7) at shoreline stations Jan-May 2005.....	39

Table 3-3.	Summary of the 2005-2006 Santa Monica Bay water quality monitoring date; surveys, dates, oceanic conditions, and directions of movement and depth of the wastewater field	40
Table 3-4.	Fecal coliform (CFU/100ml) and ammonia (mg/L) in Santa Monica Bay, Feb 2005 - Aug 2005.....	48
Table 3-5.	Fecal coliform (MPN/100ml), total coliform (MPN/100ml), enterococcus (MPN/100ml), and ammonia (mg/L) in Santa Monica Bay, Nov 2005 through Nov 2006	58
Table 3-6.	Inshore indicator bacteria percent compliance from January 2005 to May 2005.....	60
Table 3-7.	Summary of bacteriological standards for recreational waters from various sources. Values listed are for CFU/100mL or MPN/100mL	62
Table 3-8.	Inshore indicator bacteria percent compliance May 2005-December 2006..	63
Table 4-1a.	Grain size distribution of sediments in Santa Monica Bay during summer 2005.....	72
Table 4-1b.	Grain size distribution of sediments in Santa Monica Bay during summer 2006.....	73
Table 4-2a.	Pollutants in sediments of Santa Monica Bay, summer 2005. Concentrations normalized against fine grain.....	76
Table 4-2b.	Pollutants in sediments of Santa Monica Bay, summer 2006. Concentrations normalized against fine grain.....	77
Table 4-3.	Contaminant levels at outfall and reference sites. Concentrations (mg/kg for metals: μ g/kg for organics) normalized against sediment fine grains.....	90
Table 4-4a.	Concentration of metals and organic pollutants detected in sediments of Santa Monica Bay, summer 2005, with comparison to Effective Levels* (ER-L and ER-M). Concentrations in dry weight. Unit = mg/kg.....	91
Table 4-4b.	Concentration of metals and organic pollutants detected in sediments of Santa Monica Bay, summer 2006, with comparison to Effective Levels* (ER-L and ER-M). Concentrations in dry weight. Unit = mg/kg.....	92
Table 5-1.	Phylodiversity, traditional diversity indices, and community parameters for Santa Monica bay macrofauna stations, 2005-2006.....	103
Table 5-2.	Phylodiversity, traditional diversity indices, and community parameters for 7-Mile Outfall Station E6 through time.....	122

Table 5-3.	Phylodiversity, traditional diversity indices, and community parameters for 5-Mile Outfall Stations Z1 and Z2 through time.	126
Table 5-4.	Correlation table from independent contrasts of BRI and Delta against mass emission variables.	133
Table 6-1.	Taxa richness, taxonomic hierarchy composition, and taxonomic weight distribution assigned to depth category data sets based on depth category master species lists.	149
Table 6-2.	Community parameters and phylodiversity indices for Santa Monica Bay trawls, 2005-2006.	152
Table 6-3.	Pearson's product-moment correlation coefficient <i>r</i> -values of traditional community parameters and phylodiversity indices (below diagonal). Where <i>r</i> -values are significant (grey), P-values are given above diagonal in appropriate cell.	154
Table 6-4.	Community parameters and phylodiversity indices for 5-Mile Outfall trawls, 1983-2006.	165
Table 7-1a.	Trawling effort for hornyhead turbot tissue analysis during 2005 and 2006.	177
Table 7-1b.	Location and species of sport fish collected for tissue analyses during 2005 and 2006.	178
Table 7-2a.	Metals, DDT, PCB and their derivatives found in muscle and liver tissue of hornyhead turbot in Santa Monica Bay during 2005. Bolded numbers indicate DDT, PCB, and mercury levels that exceed the 0.560, 0.020, and 0.080 mg/kg OEHHA Screening Values (SV).	179
Table 7-2b.	Metals, DDT, PCB and their derivatives found in muscle and liver tissues of hornyhead turbot in Santa Monica Bay during 2006. Bold numbers indicate DDT, PCB, and mercury levels that exceed the 0.560, 0.020, and 0.080 mg/kg EPA/OEHHA SV.	181
Table 7-3.	Arsenic, selenium, mercury, and cyanide in muscle tissue of sportfish from Santa Monica Bay during 2005. Units in mg/kg wet weight.	197
Table 7-4.	Organic pollutants measured in sportfish from Santa Monica Bay during 2005. Bold numbers indicate DDT and PCB levels that exceed the 0.560 and 0.020 mg/kg SV set by EPA/OEHHA.	200

Table 7-5a. Correlation of arsenic and mercury concentrations (mg/kg) in muscle tissues of hornyhead turbot and in sediment during 2005 Santa Monica Bay monitoring 201

Table 7-5b. Correlation of arsenic and mercury concentrations (mg/kg) in muscle tissues of hornyhead turbot and in sediment during 2006 Santa Monica Bay monitoring 202