

TABLE OF CONTENTS

	Page
List of Figures.....	ii
List of Tables.....	x
Executive Summary.....	xv
Chapter 1. Introduction.....	1
Chapter 2. Effluent Quality.....	5
Chapter 3. Microbiology Water Quality - Shoreline and Inshore.....	25
Chapter 4. Water Quality - Offshore.....	45
Chapter 5. Sediment Chemistry.....	67
Chapter 6. Macrofaunal Assemblages.....	93
Chapter 7. Trawl-Caught Fish and Invertebrates.....	135
Chapter 8. Tissue Chemistry.....	167
Appendix A. Acknowledgements.....	A1
Appendix B. Diversity and Multivariate Analyses.....	B1
Appendix C. Chemical Methodologies.....	C1
Appendix D. List of Priority Pollutants Analyzed.....	D1

LIST OF FIGURES

Figure 2-1. Average Concentrations of total suspended solids, biochemical oxygen demand, and oil and grease in 5-Mile Outfall effluent from fiscal years 1986-87 through 2003-2004.....	13
Figure 2-2. Concentrations of arsenic, chromium, and lead, in plant influent and 5-Mile Outfall effluent for fiscal years 1986-87 through 2003-2004.	20
Figure 2-3. Concentrations of nickel, copper, and zinc in plant influent and 5-Mile Outfall effluent for fiscal years 1986-87 through 2003-2004.....	21
Figure 3-1. Shoreline and inshore microbiology stations in Santa Monica Bay.....	28
Figure 3-2. Monthly rainfall amounts, 2003 and 2004.	30
Figure 3-3. Annual geometric means for indicator bacteria at shoreline stations in Santa Monica Bay, 2003-2004.....	31
Figure 3-4. Annual geometric means for indicator bacteria at Santa Monica Bay shoreline stations, November and December, 2003-2004	33
Figure 3-5. Monthly geometric means for indicator bacteria for Hyperion Treatment Plant effluent, 2003-2004.....	34
Figure 3-6. Annual geometric means for indicator bacteria at inshore (surface and depth) stations in Santa Monica Bay, 2003-2004.....	38
Figure 4-1. Cooperative program water quality stations sampled during 2003 and 2004 in Santa Monica Bay.	47
Figure 4-2. Calculated salinity anomaly (S_A) and computed salinity (S) values for various dilutions based on a mean salinity of 33.346 psu	49
Figure 4-3a. Temperature 2-D plots for surveys conducted in February and May 2003 of the 45-, 60-, and 100 –m transects. The gap in May resulted from loss of data at Station 3804 due to equipment failure. Scales change by survey.....	52
Figure 4-3b. Salinity 2-D plots for surveys conducted in February and May 2003 of the 45-, 60-, and 100 –m transects. The gap in May resulted from loss of data at Station 3804 due to equipment failure. Scales change by survey.	53
Figure 4-4a. Temperature 2-D plots for surveys conducted in February and May 2004 of the 45-, 60-, and 100-m transects. Scales change by survey.....	54

Figure 4-4b. Temperature 2-D plots for surveys conducted in August and November 2004 of the 45-, 60-, and 100-m transects. Scales change by survey.....	55
Figure 4-4c. Salinity 2-D plots for surveys conducted in February and May 2004 of the 45-, 60-, and 100-m transects. Scales change by survey.....	56
Figure 4-4d. Salinity 2-D plots for surveys conducted in August and November 2004 of the 45-, 60-, and 100-m transects. Scales change by survey.....	57
Figure 4-5. Chlorophyll 3-D plots for surveys conducted in February and August 2004 depicting the absence of chlorophyll in the vicinity of the wastewater plume at depth of stratification.	58
Figure 4-6. Transmissivity and CDOM 3-D plots for May 2004 depicting an area of lowered transmissivity and colored dissolved organic material above the 5-Mile Outfall.....	59
Figure 4-7. Salinity and transmissivity 3-D plots for February 2004 survey illustrating stormwater runoff from Ballona Creek as shown by lowered salinity and transmissivity values.....	60
Figure 5-1a. Sediment chemistry stations sampled in Santa Monica Bay during 2003.	68
Figure 5-1b. Sediment chemistry stations sampled in Santa Monica Bay during 2004.	69
Figure 5-2a. Percentage of (Silt + Clay) in surface sediments of Santa Monica Bay in summer 2003.....	72
Figure 5-2b. Percentage of (Silt + Clay) in surface sediments of Santa Monica Bay in summer 2004.....	72
Figure 5-3a. Concentration of TOC in surface sediments of Santa Monica Bay in summer 2003.....	73
Figure 5-3b. Concentration of TOC in surface sediments of Santa Monica Bay in summer 2004.....	73
Figure 5-4a. Concentration of metals Ag, As, and Pb in surface sediments of Santa Monica Bay in summer 2003.....	77
Figure 5-4b. Concentration of metals Ag, As, and Pb in surface sediments of Santa Monica Bay in summer 2004.....	77
Figure 5-5a. Concentration of metals Ni and Cr in surface sediments of Santa Monica Bay in summer 2003.....	78
Figure 5-5b. Concentration of metals Ni and Cr in surface sediments of Santa Monica Bay in summer 2004.....	78

Figure 5-6a. Concentration of metals Hg and Cd in surface sediments of Santa Monica Bay in summer 2003.....	79
Figure 5-6b. Concentration of metals Hg and Cd in surface sediments of Santa Monica Bay in summer 2004.....	79
Figure 5-7a. Concentration of metals Cu and Zn in surface sediments of Santa Monica Bay in summer 2003.....	80
Figure 5-7b. Concentration of metals Cu and Zn in surface sediments of Santa Monica Bay in summer 2004.....	80
Figure 5-8a. Concentration of Total PCB and Total DDT in surface sediments of Santa Monica Bay in summer 2003.....	81
Figure 5-8b. Concentration of Total PCB and Total DDT in surface sediments of Santa Monica Bay in summer 2004.....	81
Figure 5-9a. Concentration of Total Organic Halides (TOX) in surface sediments of Santa Monica Bay in summer 2003.....	82
Figure 5-9b. Concentration of Total Organic Halides (TOX) in surface sediments of Santa Monica Bay in summer summer 2004.....	82
Figure 5-10. Concentrations (mg/kg) of chromium in sediment at E6 (7-Mile), Z2 (5-Mile) and northern reference site (C1) from 2001 to 2004.....	83
Figure 5-11. Concentrations (mg/kg) of copper in sediment at E6 (7-Mile), Z2 (5-Mile) and northern reference site (C1) from 2001 to 2004	83
Figure 5-12. Concentrations (mg/kg) of cadmium in sediment at E6 (7-Mile), Z2 (5-Mile) and northern reference site (C1) from 2001 to 2004	84
Figure 5-13. Concentrations (mg/kg) of mercury in sediment at E6 (7-Mile), Z2 (5-Mile) and northern reference site (C1) from 2001 to 2004	84
Figure 5-14. Concentrations (mg/kg) of zinc in sediment at E6 (7-Mile), Z2 (5-Mile) and northern reference site (C1) from 2001 to 2004.....	85
Figure 5-15. Concentrations (mg/kg) of Total DDT in sediment at E6 (7-Mile), Z2 (5-Mile) and northern reference site (C1) from 2001 to 2004	85
Figure 5-16. % fraction of (silt + clay) in sediment at E10, C9a, Z2 and A3 from 1995 to 2004.....	87
Figure 5-17. Concentrations of Cr, Cu, and Zn against % fraction of (silt + clay) in sediment at Z2 from 1995 to 2004.	87

Figure 6-1. Macrofaunal stations in Santa Monica Bay sampled in 2003	95
Figure 6-2. Macrofaunal stations in Santa Monica Bay sampled in 2004.....	96
Figure 6-3. Mapping of Benthic Response Index (left) and total phylogenetic diversity (right) upon mirror tree of Q-mode, parsimony analysis of endemism (PAE) cladogram, 2004.....	103
Figure 6-4. Parsimony analysis of endemism (PAE) Q-mode cladogram of Santa Monica Bay Infauna Stations, 2003.	105
Figure 6-5. Parsimony analysis of endemism (PAE) Q-mode cladogram of Santa Monica Bay Infauna Stations, 2004.	106
Figure 6-6. Principal components analysis (PCA) of 2004 Santa Monica Bay Station abiotic factors. Green depicts 18 m, brown = 130 m and greater, yellow = 70 m (d1), black = approximately 60 m, blue = 33 to 42 m, and red = 60 m 5-Mile Outfall Stations (Z1 and Z2).	107
Figure 6-7. Mirror tree mapping of DDT congeners (4,4'-DDE, left; 2,4-DDE, right) upon the Santa Monica Bay infauna parsimony analysis of endemism (PAE) Q-mode cladogram, 2004.....	108
Figure 6-8. Non-metric multidimensional scaling in 2D (above) and 3D (below) of 2004 Santa Monica Bay infauna stations derived from parsimony analysis patristic distances. Two-dimensional stress value of 0.11.....	109
Figure 6-9. Parsimony analysis of endemism (PAE) Q-mode cladogram of Santa Monica Bay Outfall Stations Z1 and Z2 through time. Post full secondary treatment depicted in green.	110
Figure 6-10. Draftsman plots of various infaunal diversity indices against time (83-04, 125 events) for Santa Monica Bay Outfall Stations Z1 and Z2.	111
Figure 6-11. Mirror tree mapping of suspended solids concentrations (left) and cyanide (right) Santa Monica Bay Outfall Stations Z1 and Z2 through time parsimony analysis of endemism (PAE) Q-mode cladogram.	113
Figure 6-12. Mirror tree mapping of Benthic Response Index (BRI, left) and total phylogenetic diversity (sPHI+, right) upon the Santa Monica Bay 5-Mile Outfall Stations Z1 and Z2 parsimony analysis of endemism (PAE) Q-mode cladogram.....	116
Figure 6-13. Principal components analysis of all community indices. 3-D mirror tree mapping of Benthic Response Index (BRI, left) and total phylogenetic diversity (sPHI+, right) upon Santa Monica Bay Outfall Stations Z1 and Z2 through time parsimony analysis of endemism (PAE) Q-mode cladogram.....	117

Figure 6-14. Mirror tree mapping of <i>Parvilucina tenuisculpta</i> (left) and <i>Capitella capitata</i> Complex (right) upon the Santa Monica Bay Outfall Stations Z1 and Z2 through time, parsimony analysis of endemism (PAE) Q-mode cladogram.....	118
Figure 6-15. Mirror tree mapping of <i>Chaetozone hartmanae</i> (left) and <i>Eclysippe trilobata</i> (right) upon the Santa Monica Bay Outfall Stations Z1 and Z2 through time, parsimony analysis of endemism (PAE) Q-mode cladogram.....	119
Figure 6-16. Within-group principal components analysis (PCA) of Santa Monica Bay Outfall Stations Z1 and Z2 through time. Pre-1999 sampling events are depicted in black and 1999-2004 are depicted in green.....	120
Figure 6-17. Non-metric multi-dimensional scaling (NMDS) derived from parsimony analysis patristic distances in 2D (above) and 3D (below) of Santa Monica Bay Outfall Stations Z1 and Z2 from 1983 to 2004.....	120
Figure 6-18. Non-metric multi-dimensional scaling (NMDS) of mass emission data (suspended solids in metric tons/year) from 1983-2004. Normalized Euclidean distances were standardized, square-root transformed, and allowed 111 restarts producing a stress value of 0.05.....	121
Figure 6-19. Abundances of select taxa at Santa Monica Bay Outfall Stations Z1 and Z2 through time.....	122
Figure 6-20. Parsimony analysis of endemism (PAE) Q-mode cladogram of peracarids at Santa Monica Bay Infauna Stations Z1 and Z2 (1983-2004).....	123
Figure 6-21. Evolutionary principal components analysis (EPCA) of Pericarida at 5-Mile Outfall Stations Z1 and Z2 through time.	124
Figure 6-22. Mirror tree mapping (solid black) of Sampling Events 85 Summer (left) and 93 Winter (right) upon the parsimony analysis of co-occurring species (PACOS) R-mode cladogram of peracarids.....	126
Figure 6-24. Mirror tree mapping (solid black) of Santa Monica Bay Outfall Sampling Events Z1 01 Summer (left) and Z2 02 (right) upon the parsimony analysis of co-occurring species (PACOS) r-mode cladogram of.....	127
Figure 6-23. Funnel plot of Santa Monica Bay 5-Mile Outfall Station Z2 showing the number of taxa versus variation in taxonomic distinctness. Dotted line represents the mean variance, while the funnel represents the 95% confidence	126
Figure 7-1. Map fo Santa Monica Bay trawl stations, 2003 and 2004	134
Figure 7-2. Depth (m) and temperature (°C) profile of Z2 Summer replicate 1 from electronic “Lotek©” archival tag device embedded in trawl door	135

Figure 7-3. Parsimony analysis of co-occurring species (PACOS) cladogram 2003. Species with shorter branch lengths were less frequently found, while those with elongate branch lengths were more frequently collected. Synapomorphies are sampling events	141
Figure 7-4 Parsimony analysis of endemism (PAE) Q-mode cladogram of Santa Monica Bay stations, 2003.....	142
Figure 7-5. Non-metric multidimensional scaling (NMDS) ordination derived from parsimony analysis of endemism (PAE) patristic distances in 2D (above) and 3D (below) of Santa Monica Bay trawl stations, 2003. Two-dimensional stress value is 0.12. Relative positions and distances of spheres reflect sampling event relationships.....	144
Figure 7-6. Parsimony analysis of co-occurring species (PACOS) cladogram, 2004. Shallow-water group from Station A1 is identified by pink circle.....	146
Figure 7-7. Parsimony analysis of endemism (PAE) Q-mode cladogram of Santa Monica Bay trawl stations.....	148
Figure 7-8. Non-metric multi-dimensional scaling (NMDS) ordination derived from parsimony analysis of endemism (PAE) patristic distances in 2D (above) and 3D (below) of Santa Monica Bay trawl stations, 2004. Two-dimensional ordination stress value is 0.09. Relative position and distances of spheres reflect sampling event relationships. Sampling Events D1T represented by yellow hexagons.....	149
Figure 7-9. Funnel plots showing average taxonomic distinctness (Δ^+) and variation in taxonomic distinctness (Δ^+) against species (taxa) richness for the 18-m depth station class of Santa Monica trawl sampling events, 2003 and 2004. The “funnel” delineates the 95% simulation limit, while the dotted line represents the taxonomic mean value of the respective index, both established by the master species list.....	150
Figure 7-10. Separate Virgulariid (sea pen) abundance (above) and composite occurrence and abundance (below) through time at Santa Monica Bay Outfall Station Z2.....	151
Figure 7-11. Pleuronectiform abundance and occurrence through time at Santa Monica Bay Outfall Station Z2. Pacific sanddab abundance (a), composite abundance (b) and composite occurrence (c).....	152
Figure 7-12. Santa Monica Bay Outfall Station Z2 through time parsimony analysis of endemism (PAE) Q-mode cladogram. Major clades and outgroup represented by red letters. Post-full-secondary treatment sampling events represented by blue lines.....	156
Figure 7-13. Mirror tree mapping of longspine combfish (left) and <i>Pleurobranchaea californica</i> (right) on the Z2 parsimony analysis of endemism (PAE) Q-mode cladogram.....	158
Figure 7-14. Within-group principal component analysis (PCA) of Santa Monica Bay Outfall Station Z2 over time. Relative positions and distances of spheres reflect sampling event relationships.....	159

Figure 7-15. Non-metric multi-dimensional scaling derived from Q-mode parsimony analysis patristic distances in 2D (above) and two orientations of 3D (below) for trawl station Z2 from 1983 to 2004. Two-dimensional ordination stress value is 0.14. Relative positions and distances of spheres reflect sampling event relationships.	160
Figure 8-1. Trawl Stations in Santa Monica Bay.	168
Figure 8-2. Zinc concentration in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	173
Figure 8-3. Arsenic concentration in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	173
Figure 8-4. Copper concentrations in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	174
Figure 8-5. Lead concentrations in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	174
Figure 8-6. Mercury concentration in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	175
Figure 8-7. Concentration of 5 metals found in muscle tissues of hornyhead turbot at station C1 during 1998 through 2002 surveys.	176
Figure 8-8. Concentrations of 5 metals found in muscle tissues of hornyhead turbot at station C3 during 1998 through 2002 surveys.	176
Figure 8-9. Concentrations of 5 metals found in muscle tissues of hornyhead turbot at station C6 during 1998 through 2002 surveys.	177
Figure 8-10. Concentrations of 5 metals found in muscle tissues of hornyhead turbot at station Z2 during 1998 through 2002 surveys.	179
Figure 8-11. Concentrations of five metals found in muscle tissues of hornyhead turbot at station C9a during 1998 through 2002 surveys.	179
Figure 8-12. Concentration of total DDT in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	180
Figure 8-13. Concentrations of total PCB in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	180
Figure 8-14. Concentrations of %lipid in muscle tissues of hornyhead turbot at five stations during 1998 through 2002 surveys.	181

Figure 8-15. Zinc concentrations in liver tissues of hornyhead turbot at five stations during 1998 through 2002 surveys..	181
Figure 8-16. Arsenic concentrations in liver tissues of hornyhead turbot at five stations during 1998 through 2002 surveys..	182
Figure 8-17. Copper concentration in liver tissues of hornyhead turbot at five stations during 1998 through 2002 surveys..	182
Figure 8-18. Lead concentration in liver tissues of hornyhead turbot at five stations during 1993 through 2000 surveys.....	186

LIST OF TABLES

Table 1-1. Summary of the City of Los Angeles’ current ocean monitoring program for Santa Monica Bay, with 1998-99 modifications.	3
Table 2-1. Constituents measured in the 5-Mile effluent monitoring program.	6
Table 2-2a. Annual averages, maximum, and minimum concentrations of conventional constituents and other pollutants in the 5-Mile Outfall effluent (January 2003 through December 2003). All concentrations are reported in µg/L unless otherwise noted.	7
Table 2-2b. Annual averages, maximum, and minimum concentrations of conventional constituents and other pollutants in the 5-Mile Outfall effluent (January 2004 through December 2004). All concentrations are reported in µg/L unless otherwise noted.	9
Table 2-3. Average, maximum, and minimum monthly averages of total suspended solids (TSS), biochemical oxygen demand (BOD), oil & grease (O&G), and settleable solids in plant influent and 5-Mile Outfall effluent during January 2003 through December 2003 and January 2004 through December 2004.	14
Table 2-4a. Average, maximum, and minimum values of detected priority pollutant metals and cyanide (µg/L) in plant influent and 5-Mile Outfall effluent during January 2003 through December 2003.	15
Table 2-4b. Average, maximum, and minimum values of detected priority pollutant metals and cyanide in plant influent and 5-Mile Outfall effluent during January 2004 through December 2004.	16
Table 2-5. Concentrations (µg/L) of key metals in 5-Mile Outfall effluent and their removal efficiency for years 1986-87 through 2003-04.	18
Table 3-1. Summary of bacteriological standards for recreational waters from various sources. Values listed are CFU/100mL.	35
Table 3-2. Percent of NPDES bacteriological standards limits compliance with total coliform standards (1-5) and E. Coli** standards (6-7) at shoreline stations, 2003-2004.	36
Table 3-3. Number of visual observations of materials of sewage origin at shoreline stations, 2003-2004.	37
Table 3-4. Percent of NPDES Permit bacteriological limits compliance with total coliform standards (1-5) and fecal coliform standards (6-7) at inshore (surface and depth) stations in Santa Monica Bay, 2003-2004.	39

Table 3-5. Storm drain flow occurrences, 2003-2004.....	41
Table 4-1. NPDES receiving water limitations.	46
Table 4-2. Summary of 2003-2004 Santa Monica Bay water quality monitoring data; surveys, dates, oceanic conditions, and direction of movement and depth of wastewater field.	50
Table 4-3. Summary of 2003-2004 Santa Monica Bay wastewater field station locations, including depth, for salinity anomaly, salinity, transmissivity, and dissolved oxy.....	51
Table 4-4. Summary of salinity anomaly values presented for the highest 15 values in descending order for each survey.....	62
Table 4-5. Compilation of salinity anomaly (S_{\ominus}), fecal coliform (CFU/100mL), and ammonia (mg/mL) from stations where both fecal coliform and ammonia were detected.	63
Table 5-1a. Grain size distribution of sediments in Santa Monica Bay during summer 2003.	70
Table 5-1b. Grain size distribution of sediments in Santa Monica Bay during summer 2004.	71
Table 5-2a. Pollutants in sediments of Santa Monica Bay, summer 2003. Concentrations normalized against fine grain. Units in mg/kg unless otherwise specified.....	75
Table 5-2b. Pollutants in sediments of Santa Monica Bay, summer 2004. Concentrations normalized against fine grain. Units in mg/kg unless otherwise specified	76
Table 5-3. Contaminant levels at outfall and reference sites. Concentrations (mg/kg for metals: $\mu\text{g}/\text{kg}$ for organics) normalized against sediment fine grains	88
Table 5-4a. Concentration of metals and organic pollutants detected in sediments of Santa Monica Bay, summer 2003, with comparison to Effective Levels* (ER-L and ER-M). Concentrations in dry weight. Unit = mg/kg	88
Table 5-4b. Concentration of metals and organic pollutants detected in sediments of Santa Monica Bay, summer 2004, with comparison to Effective Levels* (ER-L and ER-M). Concentrations in dry weight. Unit = mg/kg	89
Table 6-1. Diversity indices for Santa Monica Bay infauna stations, 2003. Species richness (S), abundance (N), Margalef (d), Pielou's evenness (J'), Fish Response Index (FRI), Shannon-Wiener loge (H'), Simpson's ($1-\lambda$), taxonomic diversity (Δ), quantitative taxonomic distinctness (Δ^*), average taxonomic distinctness (Δ^+), variation in taxonomic distinctness (Λ^+), average phylogenetic diversity (Φ^+), and total phylogenetic diversity ($s\Phi^+$) are indicated.....	100
Table 6-2. Diversity indices for Santa Monica Bay infauna stations, 2004. Species richness (S), abundance (N), Margalef (d), Pielou's evenness (J'), Fish Response Index (FRI), Shannon-Wiener loge (H'), Simpson's ($1-\lambda$), taxonomic diversity (Δ), quantitative taxonomic	

distinctness (Δ^*), average taxonomic distinctness (Δ^+), variation in taxonomic distinctness (Λ^+), average phylogenetic diversity (Φ^+), and total phylogenetic diversity ($s\Phi^+$) are indicated..... 101

Table 6-3. Diversity indices for Santa Monica Bay infauna stations Z1 and Z2 through time (83-04). Species richness (S), abundance (N), Margalef (d), Pielou's evenness (J'), Fish Response Index (FRI), Shannon-Wiener loge (H'), Simpson's (1-lambda'), taxonomic diversity (Δ), quantitative taxonomic distinctness (Δ^*), average taxonomic distinctness (Δ^+), variation in taxonomic distinctness (Λ^+), average phylogenetic diversity (Φ^+), and total phylogenetic diversity ($s\Phi^+$) are indicated..... 112

Table 7-1. Diversity indices for Santa Monica Bay trawl stations, 2003. Species richness (S), abundance (N), Margalef (d), Pielou's evenness (J'), Fish Response Index (FRI), Shannon-Wiener loge (H'), Simpson's (1-lambda'), taxonomic diversity (Δ), quantitative taxonomic distinctness (Δ^*), average taxonomic distinctness (Δ^+), variation in taxonomic distinctness (Λ^+), average phylogenetic diversity (Φ^+), and total phylogenetic diversity ($s\Phi^+$) are indicated..... 143

Table 7-2. Diversity indices for Santa Monica Bay trawl stations, 2004. Species richness (S), abundance (N), Margalef (d), Pielou's evenness (J'), Fish Response Index (FRI), Shannon-Wiener loge (H'), Simpson's (1-lambda'), taxonomic diversity (Δ), quantitative taxonomic distinctness (Δ^*), average taxonomic distinctness (Δ^+), variation in taxonomic distinctness (Λ^+), average phylogenetic diversity (Φ^+), and total phylogenetic diversity ($s\Phi^+$) are indicated..... 145

Table 7-3. Diversity indices for Santa Monica Bay trawl Station Z2 through time (83-04). Species richness (S), abundance (N), Margalef (d), Pielou's evenness (J'), Fish Response Index (FRI), Shannon-Wiener loge (H'), Simpson's (1-lambda'), taxonomic diversity (Δ), quantitative taxonomic distinctness (Δ^*), average taxonomic distinctness (Δ^+), variation in taxonomic distinctness (Λ^+), average phylogenetic diversity (Φ^+), and total phylogenetic diversity ($s\Phi^+$) are indicated..... 153

Table 7-4. Taxa from Santa Monica Bay Outfall Station Z2 having only occurred since 1999..... 159

Table 8-1a. Locations and species of trawl fish and crab samples collected during 2001 and 2002..... 169

Table 8-1b. Locations and species of sportfish samples collected during 2001 and 2002..... 169

Table 8-2a. Metals and organics found in muscle and liver tissues of hornyhead turbot in Santa Monica Bay during the 2001 sampling year. 170

Table 8-2b. Metals and organics found in muscle and liver tissues of hornyhead turbot in Santa Monica Bay during the 2002 sampling year. 171

Table 8-3a. Metals in sportfish from Santa Monica Bay during fiscal year 2001. Samples are reported in mg/kg.....	184
Table 8-3b. Metals in sportfish from Santa Monica Bay during fiscal year 2002. Samples are reported in mg/kg wet weight.	185
Table 8-4a. Organic pollutants measured in sportfish from Santa Monica Bay during 2001. Bold numbers indicate DDT and PCB levels that exceed the 0.100 ppm limit set by OEHHA (1991).	187
Table 8-4b. Organic pollutants measured in sportfish from Santa Monica Bay during 2002. Bold numbers indicate DDT and PCB levels that exceed the 0.100 ppm limit set by OEHHA (1991).	189
Table 8-5a. Correlation of metal concentrations (mg/kg) in muscle tissues of trawl fish and in sediment during 2001 Santa Monica Bay monitoring	191
Table 8-5b. Correlation of metal concentrations (mg/kg) in muscle tissues of trawl fish and in sediment during 2002 Santa Monica Bay monitoring.	191
Table C-1. Method detection limits of metals in sediments and tissues	C4

